

Gyrokinetic limit of the 2D Hartree equation in a large magnetic field

Denis Périce
dperice@constructor.university

Nicolas Rougerie
nicolas.rougerie@ens-lyon.fr

Joint work with

C>ONSTRUCTOR
UNIVERSITY

UMPA
ENS DE LYON



XLIV
Dynamics Days
EUROPE

JULY 29 - AUGUST 2, 2024
Constructor University | Bremen, Germany

Session on Mathematical and Theoretical Physics
02/08/2024

1 Context

Large system of spinless, non relativistic fermions in \mathbb{R}^2

- Homogeneous transverse magnetic field
- External potential $V : \mathbb{R}^2 \rightarrow \mathbb{R}$
- Radial interaction potential $w : \mathbb{R}^2 \rightarrow \mathbb{R}$



Motivation: Quantum Hall effect

Magnetic length: $l_b := \sqrt{\frac{\hbar}{b}}$
 \hbar : reduced Planck's constant

Semi-classical/high magnetic field limit: $l_b \rightarrow 0$

Free ground state density on \mathbb{R}^2 :

Goal: Effective dynamics



2 Model

Magnetic Laplacian

Unit system where $m = \frac{1}{2}$, $c = 1$, $q = 1$,

$$\mathcal{L}_b := (-i\hbar\nabla - b\mathbf{A})^2 = \sum_{n \in \mathbb{N}} 2\hbar b \left(n + \frac{1}{2} \right) \underbrace{\Pi_n}_{\text{projection on the } n^{\text{th}} \text{ Landau level}} \quad (1)$$

Vector potential in symmetric gauge:

$$\mathbf{A} := \frac{\mathbf{X}^\perp}{2} \implies \nabla \wedge \mathbf{A} = (\partial_1, \partial_2, \partial_3) \wedge (A_1, A_2, 0) = (0, 0, 1) \quad (2)$$

where $\mathbf{X} := (x_1, x_2)$ is the position operator.

Fermionic Density Matrix (FDM): $\gamma \in \mathcal{L}^1(L^2(\mathbb{R}^2))$ such that $\text{Tr}(\gamma) = 1$, $0 \leq \gamma \leq \frac{1}{N}$

Physical density: $\rho_\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}_+$
 $x \mapsto \gamma(x, x)$

Hartree equation:

$$i\hbar^2 \partial_t \gamma = [\mathcal{L}_b + V + w \star \rho_\gamma, \gamma] \quad (H)$$

Time scale: $l_b^{-2} = \frac{b}{\hbar}$

Scaling: $l_b \rightarrow 0$, $\hbar b = \mathcal{O}(1)$, $N = \mathcal{O}(l_b^{-2})$

Drift equation: Given a density $\rho : \mathbb{R}_+ \times \mathbb{R}^2 \rightarrow \mathbb{R}_+$,

$$\partial_t \rho(t, z) + \nabla^\perp (V + w \star \rho(t))(z) \cdot \nabla_z \rho(t, z) = 0 \quad (D)$$

3 Main result

$\Gamma(\mu, \nu)$: set of couplings between probabilities $\mu, \nu \in \mathcal{P}(\mathbb{R}^2)$, 1-Wasserstein metric:

$$W_1(\mu, \nu) := \inf_{\pi \in \Gamma(\mu, \nu)} \int_{\mathbb{R}^2 \times \mathbb{R}^2} |x - y| d\pi(x, y) \quad (3)$$

Theorem: Convergence of densities

Let γ be the solution of (H) given $\gamma(0)$ a FDM such that for some $p > 7$,

$$\text{Tr} \left(\gamma(0) \left(\mathcal{L}_b + V + \frac{1}{2} w * \rho_{\gamma(0)} \right) \right) \leq C, \quad \text{Tr}(\gamma(0) |X|^p) \leq C \quad (4)$$

Let ρ solve (D). Assume $V, w \in W^{4, \infty}(\mathbb{R}^2)$ and $\nabla w \in L^1(\mathbb{R}^2)$, $w \in H^2(\mathbb{R}^2)$.

Then, $\forall t \in \mathbb{R}_+$, $\forall \varphi \in W^{1, \infty}(\mathbb{R}^2) \cap H^2(\mathbb{R}^2)$,

$$\left| \int_{\mathbb{R}^2} \varphi(\rho_{\gamma(t)} - \rho(t)) \right| \leq \tilde{C}(t) (\|\varphi\|_{W^{1, \infty}} + \|\nabla \varphi\|_{L^2}) \left(W_1(\rho_{\gamma(0)}, \rho(0)) + I_b^{\min(2\frac{p-7}{4p-7}, \frac{2}{7})} \right) \quad (5)$$

Recap

$$i\hbar^2 \partial_t \gamma = [\mathcal{L}_b + V + w * \rho_\gamma, \gamma] \quad (H)$$

$$\text{FDM: } \gamma \in \mathcal{L}^1(L^2(\mathbb{R}^2)), \text{Tr}(\gamma) = 1, 0 \leq \gamma \leq \frac{1}{N}, \rho_\gamma(x) = \gamma(x, x)$$


$$\text{Scaling: } l_b \rightarrow 0, \quad \hbar b = \mathcal{O}(1), \quad N = \mathcal{O}(l_b^{-2})$$

$$\partial_t \rho(t, z) + \nabla^\perp (V + w * \rho(t))(z) \cdot \nabla_z \rho(t, z) = 0 \quad (D)$$

Challenges to overcome:

- Semi-classical phase space: $\mathbb{R}^2 \times \mathbb{N}$
- Controlling fast cyclotron motion
- Larger time scale

4 References



[PériceRougerie24]

D.Périce N.Rougerie. "Gyrokinetic limit of the 2D Hartree equation in a large magnetic field". In: (2024). DOI: <https://arxiv.org/abs/2403.19226>.

[Porat22]

I.B.Porat. "Derivation of Euler's equations of perfect fluids from von Neumann's equation with magnetic field". In: (2022). DOI: <https://doi.org/10.48550/arXiv.2208.01158>.

[ChenLeeLiew21]

M.Liew L.Chen J.Lee. "Combined mean-field and semiclassical limits of large fermionic systems". In: *J Stat Phys* (2021). DOI: <https://doi.org/10.1007/s10955-021-02700-w>.

[LaflecheSaffirio20]

L.Lafleche C.Saffirio. "Strong semiclassical limit from Hartree and Hartree-Fock to Vlasov-Poisson equation". In: *arXiv: Mathematical Physics* (2020). DOI: <https://arxiv.org/abs/2003.02926>.

[BenedikterPortaSchlein14]

N.Benedikter M.Porta B.Schlein. "Mean-Field Evolution of Fermionic Systems". In: *Communications in Mathematical Physics* (2014). DOI: <https://arxiv.org/abs/1305.2768>.

[GolseSaint-Raymond99]


F.Golse L.Saint-Raymond. "The Vlasov-Poisson System with Strong Magnetic Field". In: *Journal de Mathématiques Pures et Appliquées* (1999). DOI: [https://doi.org/10.1016/S0021-7824\(99\)00021-5](https://doi.org/10.1016/S0021-7824(99)00021-5).

5 Classical mechanics


Newton's second law with constant homogeneous force field F :

$$Z'' = F + bZ'^{\perp} \implies Z(t) = \underbrace{\frac{|Z'_c(0)|}{b} \begin{pmatrix} \cos(bt) \\ \sin(bt) \end{pmatrix}}_{\text{Cyclotron: } Z_c} + \underbrace{\frac{F^{\perp}}{b} t}_{\text{Drift: } Z_d \implies Z'_d = \frac{F^{\perp}}{b}} \quad \leftarrow \text{Drift time scale: } b \quad (6)$$

where we imposed $Z(0) = \frac{|Z'_c(0)|}{b} (1, 0)$, $Z'(0) = |Z'_c(0)| (0, 1) + \frac{F^{\perp}}{b}$



Classical trajectories for different b



Level sets of $V + w * \rho$

6 Semi-classical limit

Coherent state Let $\mathbf{z} := z_1 + iz_2 \in \mathbb{C}$, and $z := (z_1, z_2) \in \mathbb{R}^2$,

$$\varphi_{n,z}(x) := \frac{i^n}{\sqrt{2\pi n!} l_b} \left(\frac{x - z}{\sqrt{2} l_b} \right)^n e^{-\frac{|x-z|^2 - 2iz^\perp \cdot x}{4l_b^2}} \quad (7)$$

Phase space projector:

$$\Pi_{n,z} := |\varphi_{n,z}\rangle \langle \varphi_{n,z}|, \quad \Pi_z := \sum_{n \in \mathbb{N}} |\varphi_{n,z}\rangle \langle \varphi_{n,z}| \quad (8)$$

satisfies

$$\frac{1}{2\pi l_b^2} \int_{\mathbb{R}^2} \Pi_{n,z} dz = \Pi_n, \quad \Pi_z(x, y) = \frac{1}{2\pi l_b^2} e^{-\frac{|x-y|^2 - 2i(x^\perp \cdot y + 2z^\perp \cdot (x-y))}{4l_b^2}}, \quad \nabla_z^\perp \Pi_z = \frac{1}{il_b^2} [\Pi_z, X] \quad (*)$$

Let γ be a FDM

Phase space density

$$m_\gamma(n, z) := \frac{1}{2\pi l_b^2} \text{Tr}(\gamma \Pi_{n,z})$$

Semi-classical density

$$\rho_\gamma^{\text{sc}}(z) := \frac{1}{2\pi l_b^2} \text{Tr}(\gamma \Pi_z) = \sum_{n \in \mathbb{N}} m_\gamma(n, z)$$

Truncated semi-classical density

$$\rho_\gamma^{\text{sc}, \leq M}(z) := \sum_{n=0}^M m_\gamma(n, z)$$

Proposition: Convergence of $\rho_\gamma^{sc, \leq M}$

Let γ be a FDM, then $\forall \varphi \in L^\infty \cap H^1(\mathbb{R}^2)$,

$$\left| \int_{\mathbb{R}^2} \varphi \left(\rho_\gamma - \rho_\gamma^{sc, \leq M} \right) \right| \leq C(\varphi) \left(M^{-\frac{1}{2}} + \underbrace{\sqrt{M} l_b}_{\text{Characteristic length inside MLL}} \right) \sqrt{\text{Tr}(\gamma \mathcal{L}_b)} \quad (9)$$

We need $1 \ll M \ll l_b^{-2}$, higher Landau levels are controlled with the kinetic energy

$$\text{Tr}(\gamma \mathcal{L}_b) = 2\hbar b \sum_{n \in \mathbb{N}} \left(n + \frac{1}{2} \right) \int_{\mathbb{R}^2} m_\gamma(n, z) dz \quad (10)$$

Proposition: Gyrokinetic equation for the truncated semi-classical density

Let $t \in \mathbb{R}_+$, $\gamma(t)$ be a FDM, $W \in W^{4, \infty}(\mathbb{R}^2)$ and assume

$$i l_b^2 \partial_t \gamma(t) = [\mathcal{L}_b + W, \gamma(t)], \quad \text{Tr}(\gamma(t) \mathcal{L}_b) \leq C \quad (11)$$

then there exists a choice of $1 \ll M \ll l_b^{-2}$ such that $\forall \varphi \in L^1 \cap W^{1, \infty}(\mathbb{R}^2)$,

$$\int_{\mathbb{R}^2} \varphi \left(\partial_t \rho_{\gamma(t)}^{sc, \leq M} + \nabla^\perp W \cdot \nabla_z \rho_{\gamma(t)}^{sc, \leq M} \right) \xrightarrow{b \rightarrow \infty} 0 \quad (12)$$

Convergence $\rho_\gamma^{sc, \leq M} \rightarrow \rho$:

- Dobrushin-type stability estimate for the limiting equation
- Use confinement for initial data

7 Central computation

We recall the dynamics and (*)

$$i l_b^2 \partial_t \gamma = \text{Tr}(\mathcal{L}_b + W, \gamma), \quad \nabla_z^\perp \Pi_z = \frac{1}{i l_b^2} [\Pi_z, X]$$

Evolution part

$$\begin{aligned} \partial_t \rho_\gamma^{\text{sc}}(z) &= \frac{1}{2\pi l_b^2} \text{Tr}(\Pi_z \partial_t \gamma) = \frac{1}{2\pi l_b^2} \cdot \frac{1}{i l_b^2} \text{Tr}(\Pi_z [\mathcal{L}_b + W, \gamma]) = \frac{1}{2i\pi l_b^4} \text{Tr}(\gamma [\Pi_z, \mathcal{L}_b + W]) \\ &= \frac{1}{2i\pi l_b^4} \text{Tr}(\gamma [\Pi_z, W]) \end{aligned} \quad (13)$$

Spacial part

$$\begin{aligned} \nabla^\perp W(z) \cdot \nabla \rho_\gamma^{\text{sc}}(z) &= -\nabla W(z) \cdot \frac{1}{2\pi l_b^2} \text{Tr}(\gamma \nabla_z^\perp \Pi_z) = -\frac{1}{2i\pi l_b^4} \nabla W(z) \cdot \text{Tr}(\gamma [\Pi_z, X]) \\ &= -\frac{1}{2i\pi l_b^4} \text{Tr}(\gamma [\Pi_z, \nabla W(z) \cdot X]) \end{aligned} \quad (14)$$

so

$$\partial_t \rho_\gamma^{\text{sc}}(z) + \nabla^\perp W(z) \cdot \nabla \rho_\gamma^{\text{sc}}(z) = \frac{1}{2i\pi l_b^4} \text{Tr}(\gamma [\Pi_z, W - \nabla W(z) \cdot X]) \quad (15)$$

where

$$[\Pi_z, W - \nabla W(z) \cdot X](x, y) = \Pi_z(x, y) (W(y) - W(x) - \nabla W(z) \cdot (y - x)) \quad (16)$$

Thanks for your attention

